

INVESTIGATIONS OF THE MOST ACTIVE COMPOUND AS ANTI-DENTAL CARIES BETWEEN CHEMICAL CONSTITUENTS OF MISWAK

SALAH A JASSIM* and NABAA B ALI**

*College of Dentistry, Al Iraqia University, Baghdad-Iraq

**Dept. of Chemical Engineering, College of Engineering, University of Diayla–Diyala-Iraq

(Accepted for Publication: February 8, 2022)

ABSTRACT

The present study deals with quantum calculations to investigate some chemical descriptors of the main components of miswak extract to evaluate the most potent agents that used as anti dental caries between all the chemical components present in this plant. The chemical descriptors such as HOMO, LUMO, Energy gap, softness, hardness and electrophilicity was used as indicators to evaluate the activity of each component. Using hyperchem-8 package, density functional theory (DFT) was used as quantum method at the basis set L631G* with B3LYP function as the exchange–correlation. It can be concluded that benzyl isothiocyanate is considered as the potent agent that prevent dental careis due to the comparison of several chemical descriptors studied by DFT method.

KEY WORDS: miswak, DFT, anti-dental caries, chemical compositions.

INTRODUCTION

The potent activity of miswak as antioxidant agent come from its ability to scavenge DHHP and ABTS radicals in addition to the synergistic effect when combines with the natural antioxidant enzymes present in the oral such as peroxidase enzyme [1]. Both *Salvadora persica* and phenolic mouth washes have the same activity as dental plaque reduction and their activity to prevent the inflammation of gingival, while 8 days using of miswak can be reduced the dental plaque 75% [2-3]. Oral hygiene and periodontal diseases of teeth prevention can be attained by using *Salvadora persica* according to its anti-cariogenic activity and availability, simpilicity and its low cost [4-6]. synergistic effect of combination of miswak and Kalonji can be used to increase the antioxidant activity[7]. Chemical and mechanical effects of miswak are considered the important reason of the potent therapeutic properties of this plant. The mechanical properties include antiplaque and anticalculus in addition to cleansing, while the chemical

properties include, antimicrobial, antioxidant, anti-inflammatory and hydrating activities [4, 8-10]










Miswak is considered as good mediated to synthesis eco-frindly silver nanoparticles (AgNPs) due to the reducing agent property, which overcome the uses of toxic chemical reducing agents used for synthesis (AgNPs) [11]. miswak can be used for oral hygiene management of orthodontic patient and to prevent tooth decay [12-13]

Benzyl-isothiocyanate is considered the main constituents' active chemical compounds of miswak, it inhibits the growth and acid streptmutans production and prevent acid increase

Chemical compositions of miswak:

Different chemical components were observed in miswak by different researchers and the main one is benzyl-isothiocyanate which is more than 70% between all the components, then the rest of the materials come in different proportions such as benzaldehyde, benzyl nitrile, and others [14-22] as shown in table-1.)

Table (1):- chemical compositions of miswak

No.	Compounds	Chemical Structure
1	Benzylisothiocyanate	
2	Benzyl nitrile	
3	Carvacrol	
4	Benzaldehyde	
5	Sabinene	
6	1,8-cineole	
7	Linalool	
8	Benzyl ester	
9	Benzylisocyanate	

Computational details

The hyperchem 8 professional was used as program package. Density functional theory (DFT) was used as molecular modeling at 631G** basis sets (large size basis sets), The use of six Gaussian primitives (631G) to each core orbital improves significantly the description of the core region. B3LYP as hybrid functional was used. All the calculations are carried out at restricted Hartree-Fock levels (RHF).

Chemical descriptors were calculated according to different equations as follow [23-30]:

The hardness (η) was computed by using the following Eq:

$$\eta = (E_{\text{LUMO}} - E_{\text{HOMO}}) / 2 \quad \dots\dots(1)$$

Electrophilicity (ω) was calculated from the Eq.:

$$\omega = (E_{\text{HOMO}} + E_{\text{LUMO}} / 2)^2 / 2\eta \quad \dots\dots(2)$$

Softness (S) was calculated by:

$$S = 1 / 2\eta \quad \dots\dots(3)$$

RESULTS AND DISCUSSION

Chemical descriptors

There is a good relationship between HOMO-LUMO energy gap of the molecule and its chemical reactivity, the low value of Egap, the high active molecule (25-27), Hard molecule means that the molecule is more resistance to polarization, so it is less active when the value is high [28], While soft molecule reveals the ability of electron transfer to another molecule and the

high value of softness means more active [29]. Electrophilicity measures the energy lowering of a molecule due to maximal electron flow between the donor and acceptor, so high value of electrophilicity reflects more active molecule [29-30]

From table-2, it can be concluded from the above consideration that benzyl isothiocyanate (which is the higher percentage in miswak and reaches more than 70%) is considered the potent active between all the other chemical

constituents of miswak, and this result is matched with several experimental studies [14, 31]. The potency of benzyl isothiocyanate come from the activity of both nitrogen and sulfur atoms which are considered as active functional groups in addition to benzyl group [32].

Benzyl isothiocyanate can inhibit the growth of *Streptococcus mutans* (which is the main source of dental caries, and has fungi static action toward *Candida albicans* [33]


Table(2):- the chemical descriptors of different compositions present in miswak using DFT/ 631G**/ B3LYP


NO.	Compounds	HOMO eV	LUMO eV	Egap eV	Hardness eV	Electrophilicity eV	Softness eV
1	Benzylisothiocyanate	-6.506	-0.666	5.84	2.920	2.201	0.342
2	Benzyl nitrile	-6.955	-0.395	6.56	3.280	2.058	0.304
3	Carvacrol	-6.753	1.619	8.372	4.186	0.787	0.238
4	Benzaldehyde	-6.494	-0.407	6.087	3.043	1.955	0.328
5	Sabinene	-5.956	0.709	6.665	3.332	1.032	0.300
6	1,8-cineole	-6.205	1.865	8.07	4.035	0.583	0.247
7	Linalool	-5.847	0.373	6.22	3.110	1.204	0.321
8	Benzylester	-6.624	-0.138	6.486	3.243	1.762	0.308
9	Benzylisocyanate	-6.768	-0.215	6.553	3.276	1.860	0.305

HOMO LUMO shapes

The shapes of homo lumo are shown Fig-1 , it can be seen that homo is localized over the molecule except benzene ring while lumo is

localized above all the molecule which indicates the electrons transferred to benzene ring due to pi-pi* transition.






Fig(1): -Benzyl isothiocyanat orbitals, HOMO (A), LUMO (B)

Molecular electrostatic potential (MEP)

To investigate the nucleophilic and electrophilic attacks of the molecule and to describe the electron distribution on the reactive sites of system, (MEP) was used for this purpose which depends on the variation of colors visualized in MEP that is ranged from red color

(as negative charge that reflects the electrophilic capacity value) to blue color as positive charge. As can be seen in fig-2 , the electrophilic potency is localized on nitrogen atom which represent the more active site toward electrostatic attacks with the receptor (34):



Fig(2):- Molecular electrostatic potential (MEP) of benzyl isothiocyanat

CONCLUSION

When there is a matching pattern in the results, and when the molecules are sequenced in being the most softness and electrophilicity and the least hardness, this means a high accuracy in determining the most effective substance, so benzylisothiocyanate can inhibit *Streptococcus mutans* which is the main reason of dental caries

REFERENCES

- Saleh A Mohamed and Jalaluddin A Khan, Antioxidant capacity of chewing stick miswak *Salvadora persica*, *BMC Complementary and Alternative Medicine* 2013, 13(40): 1-6, 2013.
- Aeeza M., Mariam A., Malik S., Basil K., Marij H., Razia A. Ahmed, Comparative Clinical Effects of *Salvadora Persica* Oral Rinse and A Phenolic Commercial Mouth Wash on Human

- Oral Health; An In vivo Randomized Trial, JPDA, 30(2), 2021
- Akhilanand C., Ranjit P., and Amit N., Miswak in oral cavity –An update, J Oral Biol Craniofac Res.,3(2): 98–101, 2013
- Fatemeh E., Efficacy of Miswak (*salvadora persica*) in preventing dental caries, HEALTH , 2(5), 499-503, (2010)
- Ahtesham A., Aijaz A., Amol D. and Nilofar N., Effects of Miswak - *Salvadora Persica* on Oral Health, Al Ameen J Med Sci; 9(4), 215-218, 2016
- Mohamed M., Mathias G., Manoj K., Radha Y. and Christof D., *Salvadora persica*: Nature's Gift for Periodontal Health Antioxidants, 10(712), 1-20, 2021
- Sameen A., Kashaf J., Junaid J., Tuaha A., Waqas M., Nadia M., Kanza T., Musarrat S., Sana J. and Farheen A.i, Detection of antibacterial activities of Miswak, Kalonji and Aloe vera against oral pathogens & anti-proliferative activity against cancer cell line, BMC Complementary and Alternative Medicine, 17:265, 2-10, (2017).
- Fayez N., Mustafa N., Zohaib K. Muhammad S., Khalid A., Role of *Salvadora persica* chewing stick (miswak): A natural toothbrush for holistic oral healths, European Journal of Dentistry, 10(2): 301-308, 2016.
- Mohamed A. Faraga., Zeinab T. Shakourc, Tilo Lübkene, Andrej F., Ludger A. Wessjohannnd, Engy Mahrous, Unraveling the metabolome composition and its implication for *Salvadora persica* L. use as dental brush via a multiplex approach of NMR and LC–MS metabolomics, Journal of Pharmaceutical and Biomedical Analysis 193, 113727, 1-7, (2021)
- Haniyeh I., Gholamhoseein S., Alireza R. Dahmardeh and Zahra P. Mofrad, The Effect of Oral Care with Miswak Versus Chlorhexidine on the Incidence of Ventilator-Associated Pneumonia: A Clinical Trial Study ; 8(4):e100387, 2019
- Hammad A., Muhammad A. Sami, Saima S. & Umer H., *Salvadora persica* mediated synthesis of silver nanoparticles and their antimicrobial efficacy, Scientific Reports, 11:5996, (2021)
- Abd El-Latif H., and Sulaiman A. Alrumman, Antibacterial activity of Miswak (*Salvadora persica*) extracts against isolated and genetically identified oral cavity pathogens, Technology and Health Care 24,S841–S848, (2016).
- Khoirulzariah I., The Use of Miswak as Toothbrush for Orthodontic Patient Hindawi Publishing Corporation, Case Reports in Dentistry 2016, Article ID 7472340, 3 pages
- Abier S., Ellen M. Santangelo, Muhammad A., Anna-K. Borg-Karlson, A., Katrin P., Benzyl Isothiocyanate, a Major Component from the Roots of *Salvadora Persica* Is Highly Active against Gram-Negative Bacteria Plos One, 6(8): e23045, 2011.
- Ammar B., Guido F., Pier L. and Ivano M., The Composition of the Root Oil of *Salvadora persica* L., Journal of Essential Oil Research, 14(2):128-129, 2002.
- Hilal A. and Rajagopal K., Biological Activities of *Salvadora persica* L. (Meswak), Med Aromat Plants, 2(4):1-5. 2013
- Maged S., Elsadig H., Ahmed I. and Mohammed H., GC quantitative analysis of benzyl isothiocyanate in *Salvadora persica* roots extract and dental care herbal products, Saudi Pharm J., May; 26(4): 462–466, 2018.
- Alali F., Hudaib M., Aburjai T., Khairallah K. & Al-Hadidi N., GC-MS Analysis and Antimicrobial Activity of the Essential Oil from the Stem of the Jordanian Toothbrush Tree *Salvadora persica*. Pharmaceutical Biology, 42(8): 577–580, 2004.
- Feras A., Taha A., GC-MS Analysis and Bioactivity Testing of the Volatile Oil from the Leaves of the Toothbrush Tree *Salvadora persica* L., Formely natural product letters, 17(3):189-194, 2003.
- Arni I. Djais, Vidya Y. Tope, Effectiveness of siwak *salvadora persica* extract to aggregatibacter actinomycetemcomitans as one of pathogenic bacteria causing periodontal disease, Journal of Dentomaxillofacial Science, 2(1): 28-31, 2017.
- Saleh A. and Kasi M., Anti-biofilm activity of *Salvadora persica* on cariogenic isolates of *Streptococcus mutans*: in vitro and molecular docking studies, Biofouling, 28(1), 29-38, 2012.
- Reham A., Muhammad A., Zenebech W., Tulay L., Anna K., Borg-K., Anders G., Investigations of a Possible Chemical Effect

- of *Salvadora persica* Chewing Sticks, Evid. Based Complement., Alternat. Med. <https://doi.org/10.1155/2017/2576548>.
- Abdullah G. Al-Sehemi, Ahmad I., Sulaiman A. Alrumman4 and Abd El-Latif H., Antibacterial Activities, DFT and QSAR Studies of Quinazolinone Compounds, Bull. Chem. Soc. Ethiop., 30(2):307-316, 2016.
- Amir F. Dawood AL-Niimi , Salah A. Jassim Humadi and Naba B. Ali, Photo Electro Chemical performance evaluation of some natural dyes used in solar cells, DFT study and TD-DFT, Diyala journal for pure science, 13(3):240-248, 2017.
- Salah A. Jassim Humadi, Faliah Hassan Ali Al-Jeboori, Kafa Khalaf Hammud, Thaera A. Mussa, Synthesis and Characterization of Cu(I)-Folic Acid Complex, A Theoretical and Experimental Study, Baghdad Science Journal, 13 (2s(Supplement)) 2016.
- Salah Aldin Jassim, Evaluation of Ligand Competitive Ability for Binding in Mixed Ligand Complex, Journal of Al-Nahrain University, 19 (4): 32-40, 2016.
- Salah A. Jassim Humadi, Determination of Relative Reactivity of HSAB Complexes Using DFT Theory, Iraqi Journal of Science, 55(2B): 617-622, 2014.
- Errol G. Lewars, Computational chemistry, 2nd ed., springer, 2011.
- Zerong W., Uta W. Eusebio J., Encyclopedia of Physical Organic Chemistry, Volume 1, WILEY, 2017.
- Andrew G. Mercader, Pablo R. Duchowicz, P.M. Sivakumar, Chemometrics Applications and research: QSAR in Medicinal Chemistry, CRC press, 2016.
- Parveen D., Reet K., Luthra R., Rahul M., Gaurav S., Miswak: A periodontist's perspective, Journal of Ayurveda & Integrative Medicine, 3(4): 184-187, 2012.
- Salah A. Jassim, Removal of toxic metal ions from aqueous solutions using biosorption technique, Eurasia J Biosci 14: 3093-3099 (2020).
- Salah A. Jassim, Kafa K. Hammud, Bassam Alsheekhly, Valuable Dental Materials from *Salvadora Persica* Plants, Medico-legal Update, 21(1): 89-903, 2021
- Menna El Gaafary, Tatiana Syrovets, Hany M. Mohamed, Ahmed A. Elhenawy, Ahmed M. El-Agrody, Abd El-Galil E. Amr , Hazem A. Ghabbour and Abdulrahman A. Almehizia, Synthesis, Cytotoxic Activity, Crystal Structure, DFT Studies and Molecular Docking of 3-Amino-1-(2,5-dichlorophenyl)-8-methoxy-1H-benzo[f]chromene-2-carbonitrile, Crystals, 11(184): 1-22., 2021.